arXiv:1006.5266v1 [math.AG] 28 Jun 2010

INTEGRALITY PROPERTIES OF OPEN-CLOSED MIRROR MAPS

JIAN ZHOU

Abstract. We propose a conjecture on integrality property of the open-closed mirror maps of compact Calabi-Yau manifolds. Some examples are presented.

In this paper we will propose a conjecture on integrality properties of open-closed mirror maps for compact Calabi-Yau manifolds, inspired by the Lian-Yau integrality for closed mirror map for compact Calabi-Yau manifold [10, 11, 14, 5, 6, 7, 3], and its recent extension to the local Calabi-Yau case [12]. The integrality of open-closed mirror map in the noncompact Calabi-Yau 3-fold case was observed in [12], in this work we propose to extend it to the compact Calabi-Yau case and more noncompact examples, not necessarily three dimensional, based on a related work [13].

1. Integrality Conjecture of Open-Closed Mirror Maps

1.1. **The conjecture.** Suppose that we have a system of Picard-Fuchs equations in variables $x_0, x_1, \dots, x_N$ that comes from some charge vectors describing some brane geometry in some Calabi-Yau geometry, in the large volume phase. Then it is expected that the system has a holomorphic solution $g_0(x_0, x_1, \dots, x_N)$ such that $g_0(0, \dots, 0) = 1$ and N logarithmic solutions of the form

$$g_1^{(i)}(x_0, \dots, x_N) = \log x_i \cdot g_0(x_0, \dots, x_N) + h_1^{(i)}(x_0, \dots, x_N), \qquad i = 0, 1, \dots, N,$$

where $h_1^{(i)}$ are holomorphic functions such that $h_1^{(i)}(0, \dots, 0) = 0$. The open-closed mirror map is defined by:

$$q_i = \exp(g_1^{(i)}/g_0) = x_i \exp(h_1^{(i)}/g_0), \qquad i = 0, 1, \dots, N. \tag{1}$$

(For some examples, see e.g. [8].) On can find the inverse map by Lagrange-Good inversion formula [4] as in [12].

Our conjecture is

Conjecture 1. *The Taylor series of* $q_0, q_1, \dots, q_N$ *in* $x_0, x_1, \dots, x_N$ *are integers, and vice versa.*

We will present some examples for which we verify this conjecture. We will focus on some cases with one closed string moduli parameter. Extensions to the case of multiple closed string moduli parameters will be left to subsequent work.

1.2. **Some Picard-Fuchs systems related to open-closed mirror map.** Let $k_1, \dots, k_n$ be positive integers such that

$$\frac{1}{k_1} + \cdots + \frac{1}{k_n} = 1. \tag{2}$$

See [13] for a list of solutions for $n = 2, 3, 4, 5$ under the assumption that $k_1 \leq \cdots \leq k_n$. Here we do not require this assumption. Let $k = l.c.m.(k_1, \dots, k_n)$ and $w_i = \frac{k}{k_i}$. Then we have

$$k_1 w_1 = \cdots = k_n w_n = w_1 + \cdots + w_n = k. \tag{3}$$

Consider charge vectors formed from such a solution:

$$\begin{pmatrix} \tilde{l}^{(1)} \\ \tilde{l}^{(0)} \end{pmatrix} = \left(\begin{array}{c|cccc|cc} -k & w_1 & w_2 & \cdots & w_n & 0 & 0 \\ -1 & 1 & 0 & \cdots & 0 & -1 & 1 \end{array} \right). \tag{4}$$

We will transform them to

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|cccc|cc} -k + w_1 & 0 & w_2 & \cdots & w_n & w_1 & -w_1 \\ -1 & 1 & 0 & \cdots & 0 & -1 & 1 \end{array} \right), \tag{5}$$

where $l^{(1)} = \tilde{l}^{(1)} - w_1 \tilde{l}^{(0)}$ and $l^{(0)} = \tilde{l}^{(0)}$. We will consider the extended Picard-Fuchs system [8] associated with three charge vectors $l^{(1)}$, $l^{(0)}$ and $\tilde{l}_1 = l^{(1)} + w_1 l^{(0)}$:

$$\mathcal{L}_1 S(x_0, x_1) = 0, \qquad \mathcal{L}_0 S(x_0, x_1) = 0, \qquad \mathcal{L}_1' S(x_0, x_1) = 0, \tag{6}$$

where

$$\mathcal{L}_0 = \big[\theta_0 - x_0(1 + \theta_0 + (k - w_1)\theta_1)\big](\theta_0 - w_1\theta_1), \tag{7}$$

$$\begin{aligned} \mathcal{L}_1 = & \prod_{i=2}^{n} \prod_{j=0}^{w_i - 1} (w_i\theta_1 - j) \prod_{j=0}^{w_1 - 1} (-\theta_0 + w_1\theta_1 - j) \\ & - x_1 \prod_{j=1}^{k - w_1} (\theta_0 + (k - w_1)\theta_1 + j) \prod_{j=0}^{w_1 - 1} (-\theta_0 + w_1\theta_1 + j), \end{aligned} \tag{8}$$

$$\mathcal{L}_1' = \prod_{j=0}^{w_1 - 1} (\theta_0 - j) \cdot \prod_{i=2}^{n} \prod_{j=0}^{w_i - 1} (w_i\theta_1 - j) - x_0^{w_1} x_1 \prod_{j=1}^{k} (\theta_0 + (k - w_1)\theta_1 + j), \tag{9}$$

where $\theta_i = x_i \frac{\partial}{\partial x_i}$, $i = 0, 1$. When $n = 4$, some examples of this system correspond to the extended Picard-Fuchs system associated with B-branes in compact Calabi-Yau 3-folds in the large volume phase (see e.g. [1]). See [9] and its references for some recent work on extension of mirror symmetry to the open string case.

1.3. **Derivation of the Picard-Fuchs operators.**

Let us recall the well-known procedure of derivation of Picard-Fuchs operators from the charge vectors for the convenience of the reader. Given a vector $l = (l_0, l_1, \dots, l_{n+2}) \in \mathbb{Z}^{n+3}$, consider the affine complex space $\mathbb{C}^{p+1}$ with linear coordinates $(a_0, \dots, a_{n+2})$, define an operator

$$\mathcal{D}_l = \prod_{l_j > 0} \big(\frac{\partial}{\partial a_i}\big)^{l_j} - \prod_{l_j < 0} \big(\frac{\partial}{\partial a_i}\big)^{-l_j}. \tag{10}$$

Introduce variables

$$x_0 = \frac{a_1 a_{n+2}}{a_0 a_{n+1}}, \qquad x_1 = (-1)^k \frac{a_2^{w_2} \cdots a_n^{w_n} a_{n+1}^{w_1}}{a_0^{k - w_1} a_{n+2}^{w_1}}. \tag{11}$$

Suppose that we have a function $\tilde{\Pi}(x_0, x_1)$ such that

$$\Pi(a_0, \dots, a_{n+2}) = \frac{1}{a_0} \tilde{\Pi}(x_0, x_1) \tag{12}$$

satisfies the equations

$$\mathcal{D}_{l^{(i)}}\Pi(a_0,\dots,a_{n+2})=0,\quad i=0,1,\qquad \mathcal{D}_{\tilde{l}^{(1)}}\Pi(a_0,\dots,a_{n+2})=0. \tag{13}$$

For $i=0$,

$$\begin{aligned}\prod_{l_j^{(0)}>0}(\frac{\partial}{\partial a_i})^{l_j^{(0)}}\Pi(a_0,\dots,a_{n+2}) &= \frac{\partial}{\partial a_1}\frac{\partial}{\partial a_{n+2}}(\frac{1}{a_0}\tilde{\Pi}(x_0,x_1))\\ =\quad & \frac{1}{a_0}\frac{\partial}{\partial a_1}(\frac{1}{x_0}\frac{\partial x_0}{\partial a_{n+2}}\theta_0\tilde{\Pi}(x_0,x_1)+\frac{1}{x_1}\frac{\partial x_1}{\partial a_{n+2}}\theta_1\tilde{\Pi}(x_0,x_1))\\ =\quad & \frac{1}{a_0a_{n+2}}\frac{\partial}{\partial a_1}(\theta_0\tilde{\Pi}(x_0,x_1)-w_1\theta_1\tilde{\Pi}(x_0,x_1))\\ =\quad & \frac{1}{a_0a_{n+2}}(\frac{1}{x_0}\frac{\partial x_0}{\partial a_1}\theta_0+\frac{1}{x_1}\frac{\partial x_1}{\partial a_1}\theta_1)(\theta_0-w_1\theta_1)\tilde{\Pi}(x_0,x_1))\\ =\quad & \frac{1}{a_0a_1a_{n+2}}\theta_0(\theta_0-w_1\theta_1)\tilde{\Pi}(x_0,x_1),\end{aligned}$$

and

$$\begin{aligned}\prod_{l_j<0}(\frac{\partial}{\partial a_i})^{-l_j^{(0)}}\Pi(a_0,\dots,a_{n+2}) &= \frac{\partial}{\partial a_0}\frac{\partial}{\partial a_{n+1}}(\frac{1}{a_0}\tilde{\Pi}(x_0,x_1))\\ =\quad & \frac{\partial}{\partial a_0}(\frac{1}{a_0}\frac{1}{x_0}\frac{\partial x_0}{\partial a_{n+1}}\theta_0\tilde{\Pi}(x_0,x_1)+\frac{1}{a_0}\frac{1}{x_1}\frac{\partial x_1}{\partial a_{n+1}}\theta_1\tilde{\Pi}(x_0,x_1))\\ =\quad & \frac{1}{a_{n+1}}\frac{\partial}{\partial a_0}(\frac{1}{a_0}\cdot(-\theta_0+w_1\theta_1)\tilde{\Pi}(x_0,x_1))\\ =\quad & \frac{1}{a_{n+1}}(-\frac{1}{a_0^2}+\frac{1}{a_0}\frac{1}{x_0}\frac{\partial x_0}{\partial a_0}\theta_0+\frac{1}{a_0}\frac{1}{x_1}\frac{\partial x_1}{\partial a_0}\theta_1)(-\theta_0+w_1\theta_1)\tilde{\Pi}(x_0,x_1)\\ =\quad & \frac{1}{a_0^2a_{n+1}}(-1-\theta_0-(k-w_1)\theta_1)(-\theta_0+w_1\theta_1)\tilde{\Pi}(x_0,x_1).\end{aligned}$$

Therefore,

$$\theta_0(\theta_0-w_1\theta_1)\tilde{\Pi}(x_0,x_1)+\frac{a_1a_{n+2}}{a_0a_{n+1}}(1+\theta_0+(k-w_1)\theta_1)(-\theta_0+w_1\theta_1)\tilde{\Pi}(x_0,x_1)=0.$$

This is the equation $\mathcal{L}_0\tilde{\Pi}(x_0,x_1)=0$. The other two equations are derived in the same fashion.

1.4. **Holomorphic solutions.**

In this subsection we will show that the extended Picard-Fuch system (6) has a holomorphic solution of the form

$$g_0=1+\sum_{m_0+m_1>0}a_{m_0,m_1}x_0^{m_0}x_1^{m_1}.$$

We will derive its explicit expression as follows.

The Picard-Fuchs equations for g_0 can be rewritten as follows:

$$\begin{aligned}
& \sum_{m_0+m_1>0} m_0(m_0 - w_1m_1)a_{m_0,m_1}x_0^{m_0}x_1^{m_1} \\
= \quad & x_0 \sum_{m_0+m_1>0} (m_0 - w_1m_1)(1 + m_0 + (k - w_1)m_1)a_{m_0,m_1}x_0^{m_0}x_1^{m_1}, \\
& \sum_{m_0+m_1>0} \prod_{j=0}^{w_1-1} (-m_0 + w_1m_1 + j) \prod_{i=2}^{n} \prod_{j=0}^{w_i-1} (w_im_1 - j)a_{m_0,m_1}x_0^{m_0}x_1^{m_1} \\
= \quad & x_1 \prod_{j=1}^{k-w_1} (m_0 + (k - w_1)m_1 + j) \prod_{j=0}^{w_1-1} (-m_0 + w_1m_1 + j)a_{m_0,m_1}x_0^{m_0}x_1^{m_1}, \\
& \sum_{m_0+m_1>0} \prod_{j=0}^{w_1-1} (m_0 - j) \cdot \prod_{i=2}^{n} \prod_{j=0}^{w_i-1} (w_im_1 - j)a_{m_0,m_1}x_0^{m_0}x_1^{m_1} \\
= \quad & x_0^{w_1}x_1(k! + \sum_{m_0+m_1>0} \prod_{j=1}^{k} (m_0 + (k - w_1)m_1 + j)a_{m_0,m_1}x_0^{m_0}x_1^{m_1}).
\end{aligned}$$

These equations give rise to some recursions which can be solved directly to give us the following solution:

$$g_0(x_0, x_1) = \sum_{m\geq 0} \frac{(km)!}{\prod_{i=1}^{n}(w_im)!}(x_0^{w_1}x_1)^m. \tag{14}$$

Clearly $g_0(x_0, x_1) \in \mathbb{Z}[[x_0^{w_1}x_1]]$.

1.5. **Logarithmic solutions.** In this subsection we will show that the extended Picard-Fuch system (6) has two logarithmic solutions of the form

$$g_1^{(i)} = g_0 \log x_i + \sum_{m_0+m_1>0} a^{(i)}_{m_0,m_1}x_0^{m_0}x_1^{m_1}$$

$(i = 0, 1)$ to the Picard-Fuchs system (6). We will derive their explicit expressions as follows. From $\mathcal{L}_1' g_1^{(1)} = 0$ we get

$$\begin{aligned}
& \sum_{m_0+m_1>0} \prod_{j=0}^{w_1-1} (m_0 - j) \cdot \prod_{i=2}^{n} \prod_{j=0}^{w_i-1} (w_im_1 - j)a^{(1)}_{m_0,m_1}x_0^{m_0}x_1^{m_1} \\
+ \quad & \sum_{m=1}^{\infty} \frac{(km)!}{\prod_{i=1}^{n}(w_i(m-1))!} \sum_{i=2}^{n} \sum_{j=0}^{w_i-1} \frac{1}{w_im - j}(x_0^{w_1}x_1)^m \\
= \quad & x_0^{w_1}x_1 \sum_{m_0+m_1>0} \prod_{j=1}^{k} (m_0 + (k - w_1)m_1 + j)a^{(1)}_{m_0,m_1}x_0^{m_0}x_1^{m_1} \\
+ \quad & x_0^{w_1}x_1 \sum_{m=0}^{\infty} \frac{(k(m+1))!}{\prod_{i=1}^{n}(w_im)!} \sum_{j=1}^{k} \frac{k - w_1}{km + j}(x_0^{w_1}x_1)^m.
\end{aligned}$$

Compare the coefficients of $x_0^{m_0}x_1^{m_1}$ on both sides. When $w_0 = w_1 m$ and $w_2 = m$,

$$\begin{aligned} & a^{(1)}_{w_1 m,m} + \frac{(km)!}{\prod_{i=1}^n (w_i m)!}\sum_{i=2}^n \sum_{j=0}^{w_i-1}\frac{1}{w_i m - j} \\ = \ & \frac{\prod_{j=0}^{k-1}(km-j)}{\prod_{i=1}^n\prod_{j=0}^{w_i-1}(w_i m - j)} a^{(1)}_{w_1(m-1),m-1} + \frac{(km)!}{\prod_{i=1}^n (w_i m))!}\sum_{j=0}^{k-1}\frac{k-w_1}{km-j}. \end{aligned}$$

Hence we have

$$a^{(1)}_{w_1 m,m} = \frac{(km)!}{\prod_{i=1}^n (w_i m)!}\left(\sum_{j=1}^{km}\frac{k-w_1}{j} - \sum_{i=2}^n\sum_{j=1}^{w_i m}\frac{1}{j}\right). \tag{15}$$

For $m_1 \neq w_1 m_0$, we have

$$\begin{aligned} & \prod_{j=0}^{w_1-1}(m_0-j)\cdot\prod_{i=2}^n\prod_{j=0}^{w_i-1}(w_i m_1 - j)a^{(1)}_{m_0,m_1} \\ = \ & \prod_{j=0}^{k-1}(m_0+(k-w_1)m_1-j)a^{(1)}_{m_0-w_1,m_1-1}, \end{aligned}$$

hence when $m_0 > w_1 m_1$ we have

$$a^{(1)}_{m_0,m_1} = \frac{(m_0+(k-w_1)m_1)!}{m_0!\prod_{i=2}^n (w_i m_1)!}a^{(1)}_{m_0-w_1 m_1,0}. \tag{16}$$

We will determine $a^{(1)}_{m,0}$ from another equation.

From $\mathcal{L}_0 g_1^{(1)} = 0$ we get

$$\begin{aligned} & \sum_{m_0+m_1>0} m_0(m_0-w_1m_1)a^{(1)}_{m_0,m_1}x_0^{m_0}x_1^{m_1} - w_1\sum_{m\geq 1}\frac{(km)!w_1m}{\prod_{i=1}^n (w_i m)!}(x_0^{w_1}x_1)^m \\ = \ & x_0\sum_{m_0+m_1>0}(m_0-w_1m_1)(1+m_0+(k-w_1)m_1)a^{(1)}_{m_0,m_1}x_0^{m_0}x_1^{m_1} \\ - \ & w_1x_0\sum_{m\geq 0}\frac{(km+1)!}{\prod_{i=1}^n (w_i m)!}(x_0^{w_1}x_1)^m. \end{aligned}$$

Compare the coefficients of $x_0^{m_0}x_1^{m_1}$ on both sides. For $m_0 = m > 0$ and $m_1 = 0$,

$$a^{(1)}_{1,0} = -w_1, \qquad m^2 a^{(1)}_{m,0} = m(m-1)a^{(1)}_{m-1,0},$$

and so

$$a^{(1)}_{m,0} = -\frac{1}{m}w_1.$$

Hence by combining with (16) for $m_0 > w_1 m_1$,

$$a^{(1)}_{m_0,m_1} = -w_1\frac{(m_0+(k-w_1)m_1)!}{m_0!\prod_{i=2}^n (w_i m_1)!}\frac{1}{m_0-w_1m_1}. \tag{17}$$

When $m_0 \neq 0, w_1m_1$ or w_1m_1+1,

$$a^{(1)}_{m_0,m_1} = \frac{(m_0-1-w_1m_1)(m_0+(k-w_1)m_1)}{m_0(m_0-w_1m_1)}a^{(1)}_{m_0-1,m_1}, \tag{18}$$

therefore, for $m_0 < w_1m_1$, we have

$$a^{(1)}_{m_0,m_1} = \frac{(m_0+(k-w_1)m_1)!}{m_0!((k-w_1)m_1)!}\frac{w_1m_1}{w_1m_1-m_0}a^{(1)}_{0,m_1}. \tag{19}$$

We will determine $a^{(1)}_{0,m_1}$ from another equation.

From $\mathcal{L}_1 g^{(1)}_1 = 0$ we get:

$$\begin{aligned} & \sum_{m_0+m_1>0}\prod_{j=0}^{w_1-1}(-m_0+w_1m_1-j)\prod_{i=2}^{n}\prod_{j=0}^{w_i-1}(w_im_1-j)a^{(1)}_{m_0,m_1}x_0^{m_0}x_1^{m_1} \\ + & \sum_{m=1}^{\infty}\frac{(-1)^{w_1-1}w_1!(km)!}{(w_1m)!\prod_{i=2}^n(w_i(m-1))!}(x_0^{w_1}x_1)^m \\ = & \; x_1\sum_{m_0+m_1>0}\prod_{j=1}^{k-w_1}(m_0+(k-w_1)m_1+j)\prod_{j=0}^{w_1-1}(-m_0+w_1m_1+j)a^{(1)}_{m_0,m_1}x_0^{m_0}x_1^{m_1} \\ + & \; x_1\sum_{m=1}^{\infty}\frac{w_1!(km+k-w_1)!}{\prod_{i=1}^n(w_im)!}(x_0^{w_1}x_1)^m. \end{aligned}$$

By comparing the coefficients of $x_1^{m_1}$ we get:

$$\begin{aligned} & \prod_{i=1}^n w_i!\cdot a^{(1)}_{0,1} = w_1!(k-w_1)!, \\ & \prod_{i=1}^n\prod_{j=0}^{w_i-1}(w_im_1-j)a^{(1)}_{0,m_1} = \prod_{j=0}^{k-w_1-1}((k-w_1)m_1-j)\prod_{j=1}^{w_1}(w_1m_1-j)a^{(1)}_{0,m_1-1}, \end{aligned}$$

hence

$$a^{(1)}_{0,m_1} = \frac{((k-w_1)m_1)!}{\prod_{i=2}^n(w_im_1)!}\frac{1}{m_1}. \tag{20}$$

Combining with (19) we get:

$$a^{(1)}_{m_0,m_1} = -w_1\frac{(m_0+(k-w_1)m_1)!}{m_0!\prod_{i=2}^n(w_im)!}\frac{1}{m_0-w_1m_1}. \tag{21}$$

Therefore, we get the following solution:

$$\begin{aligned} g^{(1)}_1(x_0,x_1) = & \log x_1\cdot\sum_{m\geq 0}\frac{(km)!}{\prod_{i=1}^n(w_im)!}(x_0^{w_1}x_1)^m \\ & + \sum_{m=1}^{\infty}\frac{(km)!}{\prod_{i=1}^n(w_im)!}(\sum_{j=1}^{km}\frac{k-w_1}{j}-\sum_{i=2}^n\sum_{j=1}^{w_im}\frac{1}{j})(x_0^{w_1}x_1)^m \\ & - \sum_{m_0+m_1>0,m_0\neq w_1m_1} w_1\frac{(m_0+(k-w_1)m_1)!}{m_0!\prod_{i=2}^n(w_im)!}\frac{x_0^{m_0}x_1^{m_1}}{m_0-w_1m_1}. \end{aligned} \tag{22}$$

In the same fashion one can get the following solution:

$$\begin{aligned} g_1^{(0)}(x_0, x_1) &= \log x_0 \cdot \sum_{m\geq 0} \frac{(km)!}{\prod_{i=1}^n (w_i m)!}(x_0^{w_1} x_1)^m \\ &+ \sum_{m=1}^{\infty} \frac{(km)!}{\prod_{i=1}^n (w_i m)!}\Big(\sum_{j=1}^{km} \frac{1}{j} - \sum_{j=1}^{m} \frac{1}{w_1 j}\Big)(x_0^{w_1} x_1)^m \\ &+ \sum_{m_0+m_1>0, m_0\neq w_1 m_1} \frac{(m_0+(k-w_1)m_1)!}{m_0! \prod_{i=2}^n (w_i m)!} \frac{x_0^{m_0} x_1^{m_1}}{m_0 - w_1 m_1}. \end{aligned} \tag{23}$$

1.6. **Open-closed mirror map and its integrality.** The open-closed mirror map is given by

$$q_0 = \exp(g_1^{(0)}/g_0), \qquad q_1 = \exp(g_1^{(1)}/g_0). \tag{24}$$

As a special case of Conjecture 1, one should have

$$q_0, q_1 \in \mathbb{Z}[[x_0, x_1]]. \tag{25}$$

Notice that in both cases we have some extra terms, so the integrality property of open-closed mirror map is different from the Lian-Yau integrality of closed mirror map, because it involves the integrality of the following series:

$$\exp\Big(\sum_{m_0\neq w_1 m_1} \frac{(m_0+(k-w_1)m_1)!}{m_0! \prod_{i=2}^n (w_i m)!} \frac{x_0^{m_0} x_1^{m_1}}{m_0 - w_1 m_1} \Big/ \sum_{m\geq 0} \frac{(km)!}{\prod_{i=1}^n (w_i m)!}(x_0^{w_1} x_1)^m\Big). \tag{26}$$

Applying the Lagrange-Good inversion formula as in [12], one can get the explicit expressions for the inverse open-closed mirror map. Write

$$q_i = \sum_{m_0+m_1>0} A^{(i)}_{m_0,m_1} x_0^{m_0} x_1^{m_1}, \tag{27}$$

then $A^{(0)}_{m_0,m_1}$ and $A^{(1)}_{m_0,m_1}$ are the coefficients of $x_0^{m_0-1} x_1^{m_1}$ and $x_0^{m_0} x_1^{m_1-1}$ of

$$(1+\theta_0 h_0 + \theta_1 h_1) \cdot \exp(-m_0 h_0 - m_1 h_1), \tag{28}$$

where $h_i = g_1^{(i)}/g_0 - \log x_i$.

1.7. **The Picard-Fuchs system in a different phase.** One can also consider the Picard-Fuchs system associated with the charge vectors (4). The associated variables are:

$$\tilde{x}_0 = \frac{a_1 a_{n+2}}{a_0 a_{n+1}}, \qquad \tilde{x}_1 = (-1)^k \frac{a_1^{w_1} \cdots a_n^{w_n}}{a_0^k}. \tag{29}$$

I.e.,

$$\tilde{x}_0 = x_0, \qquad \tilde{x}_1 = x_0^{w_1} x_1. \tag{30}$$

Let $\tilde{\theta}_i = \tilde{x}_i \frac{\partial}{\partial \tilde{x}_i}$. Then we have:

$$\theta_0 = \tilde{\theta}_0 + w_1 \tilde{\theta}_1, \qquad \theta_1 = \tilde{\theta}_1, \tag{31}$$

$$\tilde{\theta}_0 = \theta_0 - w_1 \theta_1, \qquad \tilde{\theta}_1 = \theta_1. \tag{32}$$

Under such change of variables, the Picard-Fuchs operators become

$$\mathcal{L}_0 = \big[\tilde{\theta}_0 + w_1\tilde{\theta}_1 - x_0(1 + \tilde{\theta}_0 + k\tilde{\theta}_1)\big]\tilde{\theta}_0, \tag{33}$$

$$\begin{aligned}\mathcal{L}_1 = & \prod_{i=2}^{n}\prod_{j=0}^{w_i-1}(w_i\tilde{\theta}_1 - j)\prod_{j=0}^{w_1-1}(-\tilde{\theta}_0 - j) \\ & -\tilde{x}_0^{-w_1}\tilde{x}_1\prod_{j=1}^{k-w_1}(\tilde{\theta}_0 + k\tilde{\theta}_1 + j)\prod_{j=0}^{w_1-1}(-\tilde{\theta}_0 + j),\end{aligned} \tag{34}$$

$$\mathcal{L}_1' = \prod_{j=0}^{w_1-1}(\tilde{\theta}_0 + w_1\tilde{\theta}_1 - j)\cdot\prod_{i=2}^{n}\prod_{j=0}^{w_i-1}(w_i\tilde{\theta}_1 - j) - \tilde{x}_1\prod_{j=1}^{k}(\tilde{\theta}_0 + k\tilde{\theta}_1 + j). \tag{35}$$

Consider the system given by

$$\mathcal{L}_0 S(\tilde{x}_0, \tilde{x}_1) = 0, \qquad \mathcal{L}_1' S(\tilde{x}_0, \tilde{x}_1) = 0. \tag{36}$$

It has a solution holomorphic at $\tilde{x}_0 = \tilde{x}_1 = 0$:

$$\tilde{g}_0(\tilde{x}_0, \tilde{x}_1) = 1 + \sum_{m=1}^{\infty}\frac{(km)!}{\prod_{i=1}^{n}(w_i m)!}\tilde{x}_1^m, \tag{37}$$

and a logarithmic solution:

$$\tilde{g}_1^{(1)}(\tilde{x}_0, \tilde{x}_1) = \log\tilde{x}_1\cdot\tilde{g}_0(\tilde{x}_0, \tilde{x}_1) + \sum_{m=1}^{\infty}\frac{(km)!}{\prod_{i=1}^{n}(w_i m)!}\Big(\sum_{j=1}^{km}\frac{k}{j} - \sum_{i=1}^{n}\sum_{j=1}^{w_i m}\frac{1}{j}\Big)\tilde{x}_1^m. \tag{38}$$

They are not obtained from $g_0(x_0, x_1)$ and $g_1^{(1)}(x_0, x_1)$ by change of variables (30), and they do not satisfy the equation $\mathcal{L}_1 S(\tilde{x}_0, \tilde{x}_1) = 0$. Furthermore, one can easily see that there is no solution of the form $\tilde{g}_1^{(0)} = \log\tilde{x}_0\cdot\tilde{g}_0(\tilde{x}_0, \tilde{x}_1) + \tilde{h}_1^{(1)}(\tilde{x}_0, \tilde{x}_1)$, with $\tilde{h}_1^{(1)}$ holomorphic in $\tilde{x}_0$ and $\tilde{x}_1$. Indeed, if we write $\tilde{h}_1^{(0)}(\tilde{x}_0, \tilde{x}_1) = \sum_{m_0+m_1>0}\tilde{a}^{(0)}_{m_0,m_1}\tilde{x}_0^{m_0}\tilde{x}_1^{m_1}$, then from $\mathcal{L}_1'\tilde{h}_1^{(0)} = 0$ we get:

$$\begin{aligned} & \sum_{m_0+m_1>0}\prod_{j=0}^{w_1-1}(m_0 + w_1 m_1 - j)\cdot\prod_{i=2}^{n}\prod_{j=0}^{w_i-1}(w_i m_1 - j)\tilde{a}^{(0)}_{m_0,m_1}\tilde{x}_0^{m_0}\tilde{x}_1^{m_1} \\ + & \sum_{m=1}^{\infty}\sum_{j=0}^{w_1-1}\frac{1}{w_1 m - j}\cdot\frac{(km)!}{\prod_{i=1}^{n}(w_i(m-1))!}\tilde{x}_1^m \\ = & \;\tilde{x}_1\sum_{m_0+m_1>0}\prod_{j=1}^{k}(m_0 + k m_1 + j)\tilde{a}^{(0)}_{m_0,m_1}\tilde{x}_0^{m_0}\tilde{x}_1^{m_1} \\ + & \;\tilde{x}_1\sum_{m=0}^{\infty}\sum_{j=1}^{k}\frac{1}{km + j}\frac{(k(m+1))!}{\prod_{i=1}^{n}(w_i m)!}\tilde{x}_1^m.\end{aligned}$$

Compare the coefficients of $\tilde{x}_1$ on both sides:

$$0 = k!\sum_{j=1}^{k}\frac{1}{j},$$

a contradiction.

2. Integrality of Local Open-closed Mirror Maps

2.1. **Integrality of local open-closed mirror map: The outer brane case.** Consider the charge vectors

$$\begin{pmatrix}\hat{l}^{(1)}\\ \hat{l}^{(0)}\end{pmatrix}=\begin{pmatrix}\tilde{l}^{(1)}\\ -\tilde{l}^{(0)}\end{pmatrix}=\left(\begin{array}{c|cccc|cc} -k & w_1 & w_2 & \cdots & w_n & 0 & 0\\ 1 & -1 & 0 & \cdots & 0 & 1 & -1\end{array}\right). \tag{39}$$

The associated variables are

$$\hat{x}_0=\frac{a_0a_{n+1}}{a_1a_{n+2}}=\frac{1}{\tilde{x}_0},\qquad \hat{x}_1=(-1)^k\frac{a_1^{w_1}\cdots a_n^{w_n}}{a_0^k}=\tilde{x}_1. \tag{40}$$

Suppose that we have a function $\Pi(\hat{x}_0,\hat{x}_1)$ such that the following equations are satisfied:

$$\mathcal{D}_{\hat{l}^{(0)}}\Pi(\hat{x}_0,\hat{x}_1)=0,\qquad \mathcal{D}_{\hat{l}^{(1)}}\Pi(\hat{x}_0,\hat{x}_1)=0. \tag{41}$$

They give rise to the following equations:

$$\hat{\theta}_0(\hat{\theta}_0-k\hat{\theta}_1)\Pi=\hat{x}_0\hat{\theta}_0(\hat{\theta}_0-w_1\hat{\theta}_1)\Pi, \tag{42}$$

$$\prod_{j=0}^{w_1-1}(-\hat{\theta}_0+w_1\hat{\theta}_1-j)\cdot\prod_{i=2}^{n}\prod_{j=0}^{w_i-1}(w_i\hat{\theta}_1-j)\Pi=(-1)^k\hat{x}_1\prod_{j=0}^{k-1}(\hat{\theta}_0-k\hat{\theta}_1-j)\Pi. \tag{43}$$

We will refer to this system as the local extended Picard-Fuchs system associated with the charge vectors $\hat{l}^{(0)}$ and $\hat{l}^{(1)}$.

Clearly $\Phi_0=1$ is a solution, and we have the following logarithmic solutions:

$$\Phi_1^{(0)}=\log\hat{x}_0+\sum_{m\geq 1}\frac{(km)!}{\prod_{i=1}^n(w_im)!}\frac{\hat{x}_1^m}{m},\quad \Phi_1^{(1)}=\log\hat{x}_1-\sum_{m\geq 1}\frac{(km)!}{\prod_{i=1}^n(w_im)!}\frac{\hat{x}_1^m}{km}. \tag{44}$$

The local open-closed mirror map is given by

$$\hat{X}_0=\hat{x}_0\exp\sum_{m\geq 1}\frac{(km)!}{\prod_{i=1}^n(w_im)!}\frac{\hat{x}_1^m}{m},\qquad \hat{X}_1=\hat{x}_1\exp\sum_{m\geq 1}\frac{-(km)!}{\prod_{i=1}^n(w_im)!}\frac{\hat{x}_1^m}{km}. \tag{45}$$

By a result proved in [12], we have

$$\hat{x}_0^{-1}\hat{X}_0,\hat{x}_1^{-1}\hat{X}_1\in\mathbb{Z}[[\hat{x}_1]].$$

2.2. **Integrality of local open-closed mirror map: An inner brane case.** One can also consider the following charge vectors:

$$\begin{pmatrix}\tilde{\tilde{l}}^{(1)}\\ \tilde{\tilde{l}}^{(0)}\end{pmatrix}=\begin{pmatrix}\tilde{l}^{(1)}+\tilde{l}^{(0)}\\ \tilde{l}^{(0)}\end{pmatrix}=\left(\begin{array}{c|cccc|cc} -k+1 & w_1-1 & w_2 & \cdots & w_n & -1 & 1\\ -1 & 1 & 0 & \cdots & 0 & 1 & -1\end{array}\right). \tag{46}$$

The associated moduli parameters are

$$\tilde{\tilde{x}}_0=\frac{a_1a_{n+1}}{a_0a_{n+2}},\qquad \tilde{\tilde{x}}_1=(-1)^{k-1}\frac{a_1^{w_1-1}a_2^{w_2}\cdots a_n^{w_n}a_{n+2}}{a_0^{k-1}a_{n+1}}. \tag{47}$$

The local extended Picard-Fuchs system associated with these charge vectors is given by:

$$(\tilde{\hat{\theta}}_0-\tilde{\hat{\theta}}_1)(\tilde{\hat{\theta}}_0+(w_1-1)\tilde{\hat{\theta}}_1)\Pi=\tilde{\hat{y}}_0(-\tilde{\hat{\theta}}_0+\tilde{\hat{\theta}}_1)(-\tilde{\hat{\theta}}_0-(k-1)\tilde{\hat{\theta}}_1)\Pi, \tag{48}$$

$$\begin{aligned}(\tilde{\hat{\theta}}_0-\tilde{\hat{\theta}}_1)\cdot\prod_{j=0}^{w_1-2}(\tilde{\hat{\theta}}_0+(w_1-1)\tilde{\hat{\theta}}_1-j)\cdot\prod_{i=2}^{n}\prod_{j=0}^{w_i-1}(w_i\tilde{\hat{\theta}}_1-j)\Pi\\ =\hat{y}_1\cdot(-\tilde{\hat{\theta}}_0+\tilde{\hat{\theta}}_1)\prod_{j=0}^{k-2}(\tilde{\hat{\theta}}_0+(k-1)\tilde{\hat{\theta}}_1+j)\Pi.\end{aligned} \tag{49}$$

Clearly $\hat{\Phi}_0=1$ is a solution and we have two logarithmic solutions:

$$\hat{\Phi}_1^{(1)}=\log\tilde{\hat{x}}_1+\sum_{m=1}^{\infty}\frac{(km)!}{\prod_{i=1}^{n}(w_im)!}\frac{(k-1)(\tilde{\hat{x}}_0\tilde{\hat{x}}_1)^m}{km}, \tag{50}$$

$$\hat{\Phi}_1^{(0)}=\log\tilde{\hat{x}}_0-\sum_{m=1}^{\infty}\frac{(km)!}{\prod_{i=1}^{n}(w_im)!}\frac{(\tilde{\hat{x}}_0\tilde{\hat{x}}_1)^m}{km}. \tag{51}$$

The local open-closed mirror map is given by

$$\tilde{\hat{X}}_1=e^{\hat{\Phi}_1^{(1)}}=\tilde{\hat{x}}_1\exp\sum_{m=1}^{\infty}\frac{(km)!}{\prod_{i=1}^{n}(w_im)!}\frac{(k-1)(\tilde{\hat{x}}_0\tilde{\hat{x}}_1)^m}{km}, \tag{52}$$

$$\tilde{\hat{X}}_0=e^{\hat{\Phi}_1^{(0)}}=\tilde{\hat{x}}_0\exp\sum_{m=1}^{\infty}\frac{-(km)!}{\prod_{i=1}^{n}(w_im)!}\frac{(\tilde{\hat{x}}_0\tilde{\hat{x}}_1)^m}{km}. \tag{53}$$

By a result in [12], we have $\tilde{\hat{x}}_0^{-1}\tilde{\hat{X}}_0,\tilde{\hat{x}}_1^{-1}\tilde{\hat{X}}_1\in\mathbb{Z}[[\tilde{\hat{x}}_0\tilde{\hat{x}}_1]]$.

2.3. **Integrality of local open-closed mirror map: Another inner brane case.** One can also consider the the charge vectors

$$\begin{pmatrix} l^{(1)}\\ l^{(0)}\end{pmatrix}=\left(\begin{array}{c|cccc|cc} -k+w_1 & 0 & w_2 & \cdots & w_n & w_1 & -w_1\\ -1 & 1 & 0 & \cdots & 0 & -1 & 1\end{array}\right), \tag{54}$$

where $l^{(1)}=\tilde{l}^{(1)}-w_1\tilde{l}^{(0)}$ and $l^{(0)}=\tilde{l}^{(0)}$. The associated moduli parameters are

$$x_0=\frac{a_1a_{n+2}}{a_0a_{n+1}},\qquad x_1=(-1)^k\frac{a_2^{w_2}\cdots a_n^{w_n}a_{n+1}^{w_1}}{a_0^{k-w_1}a_{n+2}^{w_1}}. \tag{55}$$

The associated local extended Picard-Fuchs system is given by:

$$\check{\mathcal{L}}_1S(x_0,x_1)=0,\qquad\check{\mathcal{L}}_0S(x_0,x_1)=0,\qquad\check{\mathcal{L}}_1'S(x_0,x_1)=0, \tag{56}$$

where

$$\check{\mathcal{L}}_0=\big[\theta_0-x_0(\theta_0+(k-w_1)\theta_1)\big](\theta_0-w_1\theta_1), \tag{57}$$

$$\begin{aligned}\check{\mathcal{L}}_1=&\prod_{i=2}^{n}\prod_{j=0}^{w_i-1}(w_i\theta_1-j)\prod_{j=0}^{w_1-1}(-\theta_0+w_1\theta_1-j)\\ &-x_1\prod_{j=0}^{k-w_1-1}(\theta_0+(k-w_1)\theta_1+j)\prod_{j=0}^{w_1-1}(-\theta_0+w_1\theta_1+j),\end{aligned} \tag{58}$$

$$\check{\mathcal{L}}_1'=\prod_{j=0}^{w_1-1}(\theta_0-j)\cdot\prod_{i=2}^{n}\prod_{j=0}^{w_i-1}(w_i\theta_1-j)-x_0^{w_1}x_1\prod_{j=0}^{k-1}(\theta_0+(k-w_1)\theta_1+j). \tag{59}$$

We have a holomorphic solution $\check{g}_0 = 1$ and two logarithmic solutions:

$$\check{g}_1^{(0)} = \log x_0 + \sum_{m=1}^{\infty} \frac{(km-1)!}{\prod_{i=1}^{n}(w_i m)!}(x_0^{w_1}x_1)^m, \tag{60}$$

$$\check{g}_1^{(1)} = \log x_1 + (k-w_1)\sum_{m=1}^{\infty} \frac{(km-1)!}{\prod_{i=1}^{n}(w_i m)!}(x_0^{w_1}x_1)^m. \tag{61}$$

The local open-closed mirror map is given by

$$Q_0 = \exp(\check{g}_1^{(0)}/\check{g}_0) = x_0 \exp \sum_{m=1}^{\infty} \frac{(km-1)!}{\prod_{i=1}^{n}(w_i m)!}(x_0^{w_1}x_1)^m, \tag{62}$$

$$Q_1 = \exp(\check{g}_1^{(1)}/\check{g}_0) = x_1 \exp \sum_{m=1}^{\infty} \frac{(k-w_1)\cdot(km-1)!}{\prod_{i=1}^{n}(w_i m)!}(x_0^{w_1}x_1)^m. \tag{63}$$

By a result in [12], we have

$$x_0^{-1}Q_0, x_1^{-1}Q_1 \in \mathbb{Z}[[x_0^{w_1}x_1]]. \tag{64}$$

Using Lagrange-Good inversion formula as in [12], one can find explicit expressions for $x_0(Q_0,Q_1), x_1(Q_0,Q_1) \in \mathbb{Z}[[Q_0,Q_1]]$. Let $x_i = \sum_{m_0+m_1>0} A^{(i)}_{m_0,m_1} Q_0^{m_0}Q_1^{m_1}$. Then $A^{(0)}_{m_0,m_1}$ and $A^{(1)}_{m_0,m_1}$ are the coefficients of $x_0^{m_0-1}x_1^{m_1}$ and $x_0^{m_0}x_1^{m_1-1}$ respectively of

$$\sum_{m=1}^{\infty} \frac{(km)!}{\prod_{i=1}^{n}(w_i m)!}(x_0^{w_1}x_1)^m \cdot \exp(-m_0-(k-w_1)m_1)\sum_{m=1}^{\infty} \frac{(km-1)!}{\prod_{i=1}^{n}(w_i m)!}(x_0^{w_1}x_1)^m.$$

It follows that $x_i = Q_i \sum_{a=0}^{\infty} C_a^{(i)}(Q_0^{w_1}Q_1)^a$, where $C_a^{(0)}$ is the coefficient of y^a in

$$\sum_{m=1}^{\infty} \frac{(km)!}{\prod_{i=1}^{n}(w_i m)!}y^m \cdot \exp(-ka-1)\sum_{m=1}^{\infty} \frac{(km-1)!}{\prod_{i=1}^{n}(w_i m)!}y^m$$

and $C_a^{(1)}$ is the coefficient of y^a in

$$\sum_{m=1}^{\infty} \frac{(km)!}{\prod_{i=1}^{n}(w_i m)!}y^m \cdot \exp(-ka-(k-w_1))\sum_{m=1}^{\infty} \frac{(km-1)!}{\prod_{i=1}^{n}(w_i m)!}y^m.$$

Now we generalize the discussions in [13] to the open-closed case. Write

$$g_0(z_0,z_1) = 1 + \sum_{m_0+m_1>0} c_{m_0,m_1}q_0^{m_0}q_1^{m_1} = 1 + \sum_{m_0+m_1>0} C_{m_0,m_1}Q_0^{m_0}Q_1^{m_1}. \tag{65}$$

If Conjecture 1 holds, then the coefficients $\{c_{m_0,m_1}\}_{m_0+m_1\geq 1}$ and $\{C_{m_0,m_1}\}_{m_0+m_1\geq 1}$ are integers. One can express q_0 and q_1 as integral power series in Q_0 and Q_1, and vice versa. If Conjecture 1 holds, then the following conjecture holds.

Conjecture 2. *There are integers* $\alpha^{(i)}_{m_0,m_1}$ *and* $\beta^{(i)}_{m_0,m_1}$ *such that*

$$Q_i = q_i \prod_{m_0+m_1\geq 1}(1-q_0^{m_0}q_1^{m_1})^{\alpha^{(i)}_{m_0,m_1}}, \tag{66}$$

$$q_i = Q_i \prod_{m_0+m_1\geq 1}(1-Q_0^{m_0}Q_1^{m_1})^{\beta^{(i)}_{m_0,m_1}}. \tag{67}$$

for $i = 0, 1$.

3. Examples

We use Maple to compute some open-closed mirror maps and local open-closed mirror maps and check their integrality properties for the examples presented below.

3.1. The $n = 2$ case.

There is only one solution $k_1 = k_2 = 2$ to (2) and so $k = 2$ and $w_1 = w_2 = 1$. To the charge vectors

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|cc|cc} -1 & 0 & 1 & 1 & -1 \\ -1 & 1 & 0 & -1 & 1 \end{array} \right) \tag{68}$$

the extended Picard-Fuchs operators are:

$$\begin{aligned} \mathcal{L}_0 &= \big[\theta_0 - x_0(1 + \theta_0 + \theta_1)\big](\theta_0 - \theta_1), \\ \mathcal{L}_1 &= \theta_1(-\theta_0 + w_1\theta_1) - x_1(\theta_0 + \theta_1 + 1)(-\theta_0 + w_1\theta_1), \\ \mathcal{L}_1' &= \theta_0\theta_1 - x_0x_1 \prod_{j=1}^{2} (\theta_0 + \theta_1 + j). \end{aligned}$$

We have the following solutions:

$$\begin{aligned} g_0(x_0, x_1) &= \sum_{m \geq 0} \frac{(2m)!}{(m!)^2} (x_0x_1)^2, \\ g_1^{(0)}(x_0, x_1) &= \log x_0 \cdot \sum_{m \geq 0} \frac{(2m)!}{(m!)^2} (x_0x_1)^m + \sum_{m=1}^{\infty} \frac{(2m)!}{(m!)^2} \sum_{j=m+1}^{2m} \frac{1}{j} (x_0x_1)^m \\ &\quad + \sum_{m_0 \neq m_1} \frac{(m_0 + m_1)!}{m_0!m_1!} \frac{x_0^{m_0} x_1^{m_1}}{m_0 - m_1}, \\ g_1^{(1)}(x_0, x_1) &= \log x_1 \cdot \sum_{m \geq 0} \frac{(2m)!}{(m!)^2} (x_0x_1)^m + \sum_{m=1}^{\infty} \frac{(2m)!}{(m!)^2} \sum_{j=m+1}^{2m} \frac{1}{j} (x_0x_1)^m \\ &\quad - \sum_{m_0 \neq m_1} \frac{(m_0 + m_1)!}{m_0!m_1!} \frac{x_0^{m_0} x_1^{m_1}}{m_0 - m_1}. \end{aligned}$$

The first few terms of the Taylor series of the mirror map is given by:

$$\begin{aligned} q_0 &= x_0 + (x_0^2 - x_0x_1) + x_0^3 + (x_0^4 + x_0^3x_1 - 2x_0^2x_1^2) \\ &\quad + (x_0^5 + 2x_0^4x_1 - x_0^3x_1^2) + (x_0^6 + 3x_0^5x_1 + x_0^4x_1^2 - 5x_0^3x_1^3) + \cdots, \\ q_1 &= x_1 + (-x_0x_1 + x_1^2) + x_1^3 + (-2x_0^2x_1^2 + x_0x_1^3 + x_1^4) \\ &\quad + (2x_0x_1^4 - x_0^2x_1^3 + x_1^5) + (3x_0x_1^5 + x_0^2x_1^4 - 5x_0^3x_1^3 + x_1^6) + \cdots. \end{aligned}$$

The inverse open-closed mirror map is given by:

$$\begin{aligned} x_0 &= q_0 + (-q_0^2 + q_0q_1) + (q_0^3 - 2q_0^2q_1) + (-q_0^4 - 3q_0^2q_1^2) \\ &\quad + (q_0^5 + 2q_0^3q_1^2 + 2q_0^4q_1) + (4q_0^4q_1^2 - q_0^6 + 5q_0^3q_1^3 - 4q_0^5q_1) + \cdots, \\ x_1 &= q_1 + (-q_1^2 + q_0q_1) + (-2q_0q_1^2 + q_1^3) + (-q_1^4 - 3q_0^2q_1^2) \\ &\quad + (q_1^5 + 2q_0^2q_1^3 + 2q_0q_1^4) + (4q_0^2q_1^4 + 5q_0^3q_1^3 - q_1^6 - 4q_0q_1^5) + \cdots. \end{aligned}$$

The local extended Picard-Fuchs operators associated with (68) are:

$$\begin{aligned}\check{\mathcal{L}}_0 &= \big[\theta_0 - x_0(\theta_0+\theta_1)\big](\theta_0-\theta_1),\\ \check{\mathcal{L}}_1 &= \theta_1(-\theta_0+\theta_1) - x_1(\theta_0+\theta_1)(-\theta_0+w_1\theta_1),\\ \check{\mathcal{L}}_1' &= \theta_0\theta_1 - x_0x_1\prod_{j=0}^{1}(\theta_0+\theta_1+j).\end{aligned}$$

We have a holomorphic solution $\hat{g}_0 = 1$ and two logarithmic solutions:

$$\begin{aligned}\check{g}_1^{(0)} &= \log x_0 + \sum_{m=1}^{\infty}\frac{(2m-1)!}{(m!)^2}(x_0x_1)^m = \log\frac{2x_0}{1+\sqrt{1-4x_0x_1}},\\ \check{g}_1^{(1)} &= \log x_1 + \sum_{m=1}^{\infty}\frac{(2m-1)!}{(m!)^2}(x_0x_1)^m = \log\frac{2x_1}{1+\sqrt{1-4x_0x_1}}.\end{aligned}$$

The local open-closed mirror map is given by

$$\begin{aligned}Q_0 &= x_0\exp\sum_{m=1}^{\infty}\frac{(2m-1)!}{(m!)^2}(x_0x_1)^m = \frac{2x_0}{1+\sqrt{1-4x_0x_1}},\\ Q_1 &= x_1\exp\sum_{m=1}^{\infty}\frac{(2m-1)!}{(m!)^2}(x_0x_1)^m = \frac{2x_1}{1+\sqrt{1-4x_0x_1}}.\end{aligned}$$

One can solve these equations to get:

$$x_0 = \frac{Q_0}{1+Q_0Q_1}, \qquad x_1 = \frac{Q_1}{1-Q_0Q_1}.$$

We have checked

$$q_0 = Q_0\frac{1-Q_1}{1-Q_0}, \qquad q_1 = Q_1\frac{1-Q_0}{1-Q_1},$$

and

$$Q_0 = q_0\frac{1+q_1}{1+q_0}, \qquad Q_1 = q_1\frac{1+q_0}{1+q_1}.$$

3.2. **The $n=3$ case.** There are three cases to consider: $(k|w_1,w_2,w_3) = (3|1,1,1)$, $(4|2,1,1)$, $(6|3,2,1)$. For the first case, consider the charge vectors:

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|ccc|cc} -2 & 0 & 1 & 1 & 1 & -1 \\ -1 & 1 & 0 & 0 & -1 & 1 \end{array}\right),$$

The extended Picard-Fuchs operators are given by:

$$\begin{aligned}\mathcal{L}_0 &= \big[\theta_0 - x_0(1+\theta_0+2\theta_1)\big](\theta_0-\theta_1),\\ \mathcal{L}_1 &= \theta_1^2(-\theta_0+\theta_1) - x_1\prod_{j=1}^{2}(\theta_0+2\theta_1+j)(-\theta_0+\theta_1),\\ \mathcal{L}_1' &= \theta_0\theta_1^2 - x_0x_1\prod_{j=1}^{3}(\theta_0+2\theta_1+j),\end{aligned}$$

with solutions

$$\begin{aligned}
g_0(x_0,x_1) &= \sum_{m\geq 0}\frac{(3m)!}{(m!)^3}(x_0x_1)^m,\\
g_1^{(0)}(x_0,x_1) &= \log x_0\cdot\sum_{m\geq 0}\frac{(3m)!}{(m!)^3}(x_0x_1)^m+\sum_{m=1}^{\infty}\frac{(3m)!}{(m!)^3}\sum_{j=m+1}^{3m}\frac{1}{j}(x_0x_1)^m\\
&+ \sum_{m_0\neq m_1}\frac{(m_0+2m)!}{m_0!(m_1!)^2}\frac{x_0^{m_0}x_1^{m_1}}{m_0-m_1},\\
g_1^{(1)}(x_0,x_1) &= \log x_1\cdot\sum_{m\geq 0}\frac{(3m)!}{(m!)^3}(x_0x_1)^m+2\sum_{m=1}^{\infty}\frac{(3m)!}{(m!)^3}\sum_{j=m+1}^{3m}\frac{1}{j}(x_0x_1)^m\\
&- \sum_{m_0\neq m_1}w_1\frac{(m_0+2m_1)!}{m_0!(m!)^2}\frac{x_0^{m_0}x_1^{m_1}}{m_0-m_1}.
\end{aligned}$$

The open-closed mirror map is given by:

$$\begin{aligned}
q_0 &= x_0+(x_0^2-2x_0x_1)+(x_0^3+3x_0^2x_1-x_0x_1^2)+(x_0^4+9x_0^3x_1-29x_0^2x_1^2-2x_0x_1^3)\\
&+ (x_0^5+16x_0^4x_1+27x_0^3x_1^2-23x_0^2x_1^3-5x_0x_1^4)\\
&+ (x_0^6+24x_0^5x_1+127x_0^4x_1^2-527x_0^3x_1^3-63x_0^2x_1^4-14x_0x_1^5)+\cdots,\\
q_1 &= x_1+(-x_0x_1+2x_1^2)+(8x_0x_1^2+5x_1^3)+(-16x_0^2x_1^2+33x_0x_1^3+14x_1^4)\\
&+ (-x_0^3x_1^2+111x_0^2x_1^3+124x_0x_1^4+42x_1^5)\\
&+ (-x_0^4x_1^2-307x_0^3x_1^3+606x_0^2x_1^4+462x_0x_1^5+132x_1^6)+\cdots.
\end{aligned}$$

The local open-closed mirror map is given by:

$$Q_0=x_0\exp\sum_{m=1}^{\infty}\frac{(3m-1)!}{(m!)^3}(x_0x_1)^m,$$

$$Q_1=x_1\exp\sum_{m=1}^{\infty}\frac{2\cdot(3m-1)!}{(m!)^3}(x_0x_1)^m.$$

For $(k|w_1,w_2,w_3)=(4|2,1,1)$, one can consider the charge vectors:

$$\begin{pmatrix} l^{(1)}\\ l^{(0)}\end{pmatrix}=\left(\begin{array}{c|ccc|cc} -2 & 0 & 1 & 1 & 2 & -2\\ -1 & 1 & 0 & 0 & -1 & 1\end{array}\right),$$

The open-closed mirror map is given by:

$$\begin{aligned}
q_0 &= x_0+(x_0^2-x_0x_1)+(x_0^3-7x_0^2x_1-x_0x_1^2)+(x_0^4+12x_0^3x_1-5x_0^2x_1^2-2x_0x_1^3)\\
&+ (x_0^5+20x_0^4x_1-39x_0^3x_1^2-14x_0^2x_1^3-5x_0x_1^4)\\
&+ (x_0^6+29x_0^5x_1-299x_0^4x_1^2-80x_0^3x_1^3-45x_0^2x_1^4-14x_0x_1^5)+\cdots,\\
q_1 &= x_1+(-2x_0x_1+2x_1^2)+(x_0^2x_1+8x_0x_1^2+5x_1^3)+(4x_0^2x_1^2+34x_0x_1^3+14x_1^4)\\
&+ (-56x_0^3x_1^2+107x_0^2x_1^3+128x_0x_1^4+42x_1^5)\\
&+ (40x_0^4x_1^2+220x_0^3x_1^3+592x_0^2x_1^4+476x_0x_1^5+132x_1^6)+\cdots.
\end{aligned}$$

The inverse open-closed mirror map is given by:

$$\begin{aligned}
x_0 &= q_0+(-q_0^2+q_0q_1)+(q_0^3+6q_0^2q_1)+(-q_0^2q_1^2-q_0^4-25q_0^3q_1)+\cdots,\\
x_1 &= q_1+(2q_0q_1-2q_1^2)+(q_0^2q_1-18q_0q_1^2+3q_1^3)+(50q_0q_1^3-56q_0^2q_1^2-4q_1^4)+\cdots.
\end{aligned}$$

The local open-closed mirror map is given by:

$$Q_0 = x_0 \exp \sum_{m=1}^{\infty} \frac{(4m-1)!}{(2m)!(m!)^2} (x_0^2 x_1)^m,$$
$$Q_1 = x_1 \exp \sum_{m=1}^{\infty} \frac{2\cdot(4m-1)!}{(2m)!(m!)^2} (x_0^2 x_1)^m.$$

The inverse local open-closed mirror map is given by:

$$\begin{aligned} x_0 &= Q_0(1 - 3Q_0^2Q_1 - 12Q_0^4Q_1^2 - 253Q_0^6Q_1^3 - 6033Q_0^8Q_1^4 + \cdots), \\ x_1 &= Q_1(1 - 6Q_0^2Q_1 - 15Q_0^4Q_1^2 - 434Q_0^6Q_1^3 - 10404Q_0^8Q_1^4 + \cdots). \end{aligned}$$

One has

$$\begin{aligned} q_0 \quad = \quad & Q_0 - Q_0Q_1 + Q_0^2 - Q_0Q_1^2 - 7Q_0^2Q_1 + Q_0^3 - 2Q_0Q_1^3 - 5Q_0^2Q_1^2 + 9Q_0^3Q_1 + Q_0^4 \\ & -5Q_0Q_1^4 - 14Q_0^2Q_1^3 - 30Q_0^3Q_1^2 + 14Q_0^4Q_1 + Q_0^5 \\ & -14Q_0Q_1^5 - 45Q_0^2Q_1^4 - 65Q_0^3Q_1^3 - 215Q_0^4Q_1^2 + 20Q_0^5Q_1 + Q_0^6 + \cdots \\ q_1 \quad = \quad & Q_1 + 2Q_1^2 - 2Q_0Q_1 + 5Q_1^3 + 8Q_0Q_1^2 + Q_0^2Q_1 + 14Q_1^4 + 34Q_0Q_1^3 - 2Q_0^2Q_1^2 \\ & +42Q_1^5 + 128Q_0Q_1^4 + 83Q_0^2Q_1^3 - 38Q_0^3Q_1^2 \\ & +132Q_1^6 + 476Q_0Q_1^5 + 502Q_0^2Q_1^4 + 100Q_0^3Q_1^3 + 28Q_0^4Q_1^2 + 1776Q_0Q_1^6 + \cdots . \end{aligned}$$

One can also consider the charge vectors corresponding to another phase of the B-brane:

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|ccc|cc} -3 & 0 & 2 & 1 & 1 & -1 \\ -1 & 1 & 0 & 0 & -1 & 1 \end{array} \right),$$

Then the open-closed mirror map is given by:

$$\begin{aligned} q_0 &= x_0 + (x^2 - 3x_0x_1) + (x_0^3 + 10x_0^2x_1 - 3x_0x_1^2) \\ &+ (x^4 + 28x_0^3x_1 - 111x_0^2x_1^2 - 10x_0x_1^3) + \cdots , \\ q_1 &= x_1 + (-x_0x_1 + 3x_1^2) + (42x_0x_1^2 + 12x_1^3) \\ &+ (-63x_0^2x_1^2 + 192x_0x_1^3 + 55x_1^4) + \cdots . \end{aligned}$$

For $(k|w_1, w_2, w_3) = (6|3, 2, 1)$, we check the following three cases corresponding to three different phases of the B-brane: For the charge vectors

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|ccc|cc} -3 & 0 & 2 & 1 & 3 & -3 \\ -1 & 1 & 0 & 0 & -1 & 1 \end{array} \right),$$

we have

$$\begin{aligned} q_0 &= x_0 + (x_0^2 - x_0x_1) + (x_0^3 - 7x_0^2x_1 - 2x_0x_1) \\ &+ (x_0^4 - 37x_0^3x_1 - 17x_0^2x_1^2 - 7x_0x_1^3) + \cdots , \\ q_1 &= x_1 + (-3x_0x_1 + 3x_1^2) + (3x_0^2x_1 + 9x_0x_1^2 + 12x_1^3) \\ &+ (-x_0^3x_1 + 45x_0^2x_1^2 + 81x_0x_1^3 + 55x_1^4) + \cdots ; \end{aligned}$$

for the charge vectors

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|ccc|cc} -4 & 0 & 3 & 1 & 2 & -2 \\ -1 & 1 & 0 & 0 & -1 & 1 \end{array} \right),$$

we have

$$\begin{aligned} q_0 &= x_0 + (x_0^2 - 2x_0x_1) + (x_0^3 - 22x_0^2x_1 - 5x_0x_1^2) \\ &+ (x_0^4 + 212x_0^3x_1 - 49x_0^2x_1^2 - 24x_0x_1^3) + \cdots , \\ q_1 &= x_1 + (-2x_0x_1 + 4x_1^2) + (x_0^2x_1 + 32x_0x_1^2 + 22x_1^2) \\ &+ (342x_0^2x_1^2 + 284x_0x_1^3 + 140x_1^4) + \cdots ; \end{aligned}$$

for the charge vectors

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|ccc|cc} -5 & 0 & 2 & 3 & 1 & -1 \\ -1 & 1 & 0 & 0 & -1 & 1 \end{array} \right),$$

we have

$$\begin{aligned} q_0 &= x_0 + (x_0^2 - 10x_0x_1) + (x_0^3 + 77x_0^2x_1 - 55x_0x_1^2) \\ &+ (x_0^4 + 227x_0^3x_1 - 2635x_0^2x_1^2 - 785x_0x_1^3) + \cdots , \\ q_1 &= x_1 + (-x_0x_1 + 10x_1^2) + (525x_0x_1^2 + 155x_1^3) \\ &+ (-685x_0^2x_1^2 + 6905x_0x_1^3 + 2885x_1^4) + \cdots . \end{aligned}$$

3.3. **Some $n > 3$ cases.** For $n = 4$, there are thirteen cases to consider when $w_1 \geq \cdots \geq w_4$ listed in [13]; when $n = 5$ there are 147 cases. We have checked some of them using our Maple algorithm. The following cases are particularly interesting because they correspond to the large volume phase of B-brane geometries in one-moduli Calabi-Yau hypersurfaces in weighted projective spaces:

$$(k|w_1, \dots, w_5) = (5|1,1,1,1,1), (6|2,1,1,1,1), (8|4,1,1,1,1), (10|5,2,1,1,1).$$

When $(k|w_1, \dots, w_5) = (5|1,1,1,1,1)$, for the charge vectors

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|ccccc|cc} -4 & 0 & 1 & 1 & 1 & 1 & -1 & 1 \\ -1 & 1 & 0 & 0 & 0 & 0 & 1 & -1 \end{array} \right) \tag{69}$$

we have

$$\begin{aligned} q_0 &= x_0 + (x_0^2 - 24x_0x_1) + (x_0^3 - 24x_0^2x_1 - 972x_0x_1^2) \\ &+ (x_0^4 + 216x_0^3x_1 - 20772x_0^2x_1^2 - 95264x_0x_1^3) + \cdots , \\ q_1 &= x_1 + (-x_0x_1 + 24x_1^2) + (746x_0x_1^2 + 1548x_1^3) \\ &+ (-1010x_0^2x_1^2 + 36732x_0x_1^3 + 155744x_1^4) + \cdots . \end{aligned}$$

When $(k|w_1, \dots, w_5) = (6|2,1,1,1,1)$, we consider two phases. For the charge vectors

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|ccccc|cc} -4 & 0 & 1 & 1 & 1 & 1 & -2 & 2 \\ -1 & 1 & 0 & 0 & 0 & 0 & 1 & -1 \end{array} \right)$$

we have

$$\begin{aligned} q_0 &= x_0 + (x_0^2 - 12x_0x_1) + (x_0^3 - 132x_0^2x_1 - 558x_0x_1^2) \\ &+ (x_0^4 + 570x_0^3x_1 - 6678x_0^2x_1^2 - 54328x_0x_1^3) + \cdots , \\ q_1 &= x_1 + (-2x_0x_1 + 24x_1^2) + (x_0^2x_1 + 192x_0x_1^2 + 1548x_1^3) \\ &+ (1632x_0^2x_1^2 + 17784x_0x_1^3 + 155744x_1^4) + \cdots ; \end{aligned}$$

for the charge vectors

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|ccccc|cc} -5 & 0 & 2 & 1 & 1 & 1 & -1 & 1 \\ -1 & 1 & 0 & 0 & 0 & 0 & 1 & -1 \end{array} \right)$$

we have

$$\begin{aligned} q_0 &= x_0 + (x_0^2 - 60x_0x_1) + (x_0^3 + 462x_0^2x_1 - 7650x_0x_1^2) \\ &+ (x_0^4 + 1362x_0^3x_1 - 225270x_0^2x_1^2 - 2271800x_0x_1^3) + \cdots , \\ q_1 &= x_1 + (-x_0x_1 + 60x_1^2) + (-60x_0x_1^2 + 11250x_1^3) \\ &+ (1890x_0^2x_1^2 + 175050x_0x_1^3 + 3405800x_1^4) + \cdots . \end{aligned}$$

When $(k|w_1,\ldots,w_5) = (8|4,1,1,1,1)$, for the charge vectors

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|ccccc|cc} -4 & 0 & 1 & 1 & 1 & 1 & -4 & 4 \\ -1 & 1 & 0 & 0 & 0 & 0 & 1 & -1 \end{array} \right)$$

we have

$$\begin{aligned} q_0 &= x_0 + (x_0^2 - 6x_0x_1) + (x_0^3 - 46x_0^2x_1 - 297x_0x_1^2) \\ &+ (x_0^4 - 226x_0^3x_1 - 3297x_0^2x_1^2 - 28946x_0x_1^3) + \cdots , \\ q_1 &= x_1 + (-4x_0x_1 + 24x_1^2) + (6x_0^2x_1 + 64x_0x_1^2 + 1548x_1^3) \\ &+ (-4x_0^3x_1 + 224x_0^2x_1^2 + 10608x_0x_1^3 + 155744x_1^4) + \cdots ; \end{aligned}$$

for the charge vectors

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|ccccc|cc} -7 & 0 & 4 & 1 & 1 & 1 & -1 & 1 \\ -1 & 1 & 0 & 0 & 0 & 0 & 1 & -1 \end{array} \right) \tag{70}$$

we have

$$\begin{aligned} q_0 &= x_0 + (x_0^2 - 210x_0x_1) + (x_0^3 + 2676x_0^2x_1 - 113085x_0x_1^2) \\ &+ (x_0^4 + 8556x_0^3x_1 - 4420395x_0^2x_1^2 - 137765950x_0x_1^3) + \cdots , \\ q_1 &= x_1 + (-x_0x_1 + 210x_1^2) + (23212x_0x_1^2 + 157185x_1^3) \\ &+ (-29302x_0^2x_1^2 + 8462685x_0x_1^3 + 194522650x_1^4) + \cdots . \end{aligned}$$

When $(k|w_1,\ldots,w_5) = (10|5,2,1,1,1)$, for the charge vectors

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|ccccc|cc} -5 & 0 & 2 & 1 & 1 & 1 & -5 & 5 \\ -1 & 1 & 0 & 0 & 0 & 0 & 1 & -1 \end{array} \right)$$

we have

$$\begin{aligned} q_0 &= x_0 + (x_0^2 - 12x_0x_1) + (x_0^3 - 102x_0^2x_1 - 1818x_0x_1^2) \\ &+ (x_0^4 - 522x_0^3x_1 - 23838x_0^2x_1^2 - 538168x_0x_1^3) + \cdots , \\ q_1 &= x_1 + (-5x_0x_1 + 60x_1^2) + (10x_0^2x_1 + 150x_0x_1^2 + 11250x_1^3) \\ &+ (-10x_0^3x_1 + 450x_0^2x_1^2 + 86250x_0x_1^3 + 3405800x_1^4) + \cdots ; \end{aligned}$$

for the charge vectors

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|ccccc|cc} -8 & 0 & 5 & 1 & 1 & 1 & -2 & 2 \\ -1 & 1 & 0 & 0 & 0 & 0 & 1 & -1 \end{array} \right) \tag{71}$$

we have

$$\begin{aligned} q_0 &= x_0 + (x_0^2 - 168x_0x_1) + (x_0^3 - 3192x_0^2x_1 - 166068x_0x_1^2) \\ &+ (x_0^4 + 33534x_0^3x_1 - 3742116x_0^2x_1^2 - 336621152x_0x_1^3) + \cdots , \\ q_1 &= x_1 + (-2x_0x_1 + 336x_1^2) + (x_0^2x_1 + 5376x_0x_1^2 + 416808x_1^3) \\ &+ (-11760x_0^2x_1^2 + 9366672x_0x_1^3 + 859605376x_1^4) + \cdots ; \end{aligned}$$

for the charge vectors

$$\begin{pmatrix} l^{(1)} \\ l^{(0)} \end{pmatrix} = \left(\begin{array}{c|ccccc|cc} -9 & 0 & 5 & 2 & 1 & 1 & -1 & 1 \\ -1 & 1 & 0 & 0 & 0 & 0 & 1 & -1 \end{array} \right) \tag{72}$$

we have

$$\begin{aligned} q_0 &= x_0 + (x_0^2 - 1512 x_0 x_1) + (x_0^3 + 27654 x_0^2 x_1 - 8046108 x_0 x_1^2) \\ &+ (x_0^4 + 95694 x_0^3 x_1 - 378472500 x_0^2 x - 1^2 - 93766645728 x_0 x_1^3) + \cdots, \\ q_1 &= x_1 + (-x_0 x_1 + 1512 x_1^2) + (265332 x_0 x_1^2 + 10332252 x_1^3) \\ &+ (121554726048 x_1^4 + 719463276 x_0 x_1^3 - 334884 x_0^2 x_1^2) + \cdots. \end{aligned}$$

Some $n > 4$ cases have also been checked.

Acknowledgements. This research is partly supported by NSFC grants (10425101 and 10631050) and a 973 project grant NKBRPC (2006cB805905).

References

[1] M. Alim, M. Hecht, P. Mayr, A. Mertens *Mirror Symmetry for Toric Branes on Compact Hypersurfaces*, arXiv:0901.2937.

[2] T.-M. Chiang, A. Klemm, S.-T. Yau, E. Zaslow, *Local Mirror Symmetry: Calculations and Interpretations*, Adv.Theor.Math.Phys. **3** (1999), 495-565.

[3] E. Delaygue, *Critére pour l'intégralité des coefficients de Taylor des applications miroir*, arXiv:0912.3776.

[4] I. J. Good, *Generalizations to several variables of Lagrange's expansion, with applications to stochastic processes*, Proc. Cambridge Philos. Soc. **56** (1960), 367-380.

[5] C. Krattenthaler,T. Rivoal, *Multivariate p-adic formal congruences and integrality of Taylor coefficients of mirror maps*, arXiv:0804.3049.

[6] C. Krattenthaler, T. Rivoal, *On the integrality of the Taylor coefficients of mirror maps*, Duke Math. J. **151** (2010), 175-218, arXiv:0907.2577.

[7] C. Krattenthaler, T. Rivoal, *On the integrality of the Taylor coefficients of mirror maps, II*, Commun. Number Theory Phys. **3** (2009), 555-591, arXiv:0907.2578.

[8] W. Lerche, P. Mayr, *On $N = 1$ mirror symmetry for open type II strings*, arXiv:hep-th/0111113.

[9] S. Li, B.H. Lian, S.-T. Yau, *Picard-Fuchs equations for relative periods and Abel-Jacobi map for Calabi-Yau hypersurfaces*, arXiv:0910.4215.

[10] B. H. Lian, S.-T. Yau, *Mirror maps, modular relations and hypergeometric series I*, appeared as *Integrality of certain exponential series* , in: Lectures in Algebra and Geometry, Proceedings of the International Conference on Algebra and Geometry, Taipei, 1995, M.-C. Kang (ed.), Int. Press, Cambridge, MA, 1998, pp. 215-227.

[11] B. H. Lian, S.-T. Yau, *The nth root of the mirror map*, in: Calabi-Yau varieties and mirror symmetry, Proceedings of the Workshop on Arithmetic, Geometry and Physics around Calabi-Yau Varieties and Mirror Symmetry, Toronto, ON, 2001, N. Yui and J. D. Lewis (eds.), Fields Inst. Commun., 38, Amer. Math. Soc., Providence, RI, 2003, pp. 195-199.

[12] J. Zhou, *Some integrality properties in local mirror symmetry*, arXiv:1005.3243.

[13] J. Zhou, *Integrality properties of variations of Mahler measures*, arXiv:1006.2428.

[14] W. Zudilin, *Integrality of power expansions related to hypergeometric series*, Mathematical Notes **71.5** (2002), 604-616.

Department of Mathematical Sciences, Tsinghua University, Beijing, 100084, China
E-mail address: `jzhou@math.tsinghua.edu.cn`